

On Weakly Sign-Symmetric Matrices

B. M. GREEN*

*Oregon College of Education
Monmouth, Oregon*

Recommended by David Carlson

ABSTRACT

Weakly sign-symmetric matrices have non-negative principal minors and non-negative products of symmetrically placed pairs of almost-principal minors. A necessary condition is proved for such a matrix to have as rank a given positive integer. Several characterizations are given of those weakly sign-symmetric matrices for which the generalized Hadamard inequality holds.

INTRODUCTION

We use terminology and notation similar to that of Carlson [1], [2], and Marcus and Minc [10]. Let α, β and γ, \dots be subsets of $Z_n = \{1, 2, \dots, n\}$ and assumed to be in their natural order. Let $|\alpha|$ be the order of α . Let A be an $n \times n$ complex matrix; then $A[\alpha|\beta]$ is the submatrix of A with rows and columns indexed by α and β , respectively, and if $|\alpha| = |\beta|$, $A(\alpha|\beta) = \det A[\alpha|\beta]$. We let $A[\alpha] = A[\alpha|\alpha]$ and $A(\alpha) = A(\alpha|\alpha)$; if α is empty then $A(\alpha) = 1$. Next for convenience let i be the complement of i in Z_n , let $\alpha - i = \alpha - \{i\}$, and let $\{\alpha, i\} = \alpha \cup \{i\}$.

The term A is a weakly sign-symmetric matrix or WSS-matrix if and only if $A(\alpha) \geq 0$ for all $\alpha \subseteq Z_n$ and

$$A(\alpha, i|\beta, j)A(\alpha, j|\alpha, i) \geq 0 \quad \text{for all } \alpha \subseteq Z_n, \quad i, j \in Z_n - \alpha. \quad (1)$$

* This work was begun while the author held an NSF Science Faculty Fellowship and is part of the author's Doctoral dissertation which was directed by Professor David Carlson.

A minor of the type $A(\alpha, i|\alpha, j)$, $\alpha \subseteq Z_n$, $i, j \in Z_n - \alpha$, will be called almost principal.

The term A is a generalized Hadamard or GH-matrix if and only if $A(\alpha) \geq 0$ for all $\alpha \subseteq Z_n$ and

$$A(\alpha \cup \beta)A(\alpha \cap \beta) \leq A(\alpha)A(\beta) \quad \text{for all } \alpha, \beta \subseteq Z_n. \quad (2)$$

The classical inequalities of Hadamard

$$a_{11}a_{22} \cdots a_{nn} \geq \det A,$$

and Fischer,

$$A(\alpha \cup \beta) \leq A(\alpha)A(\beta) \quad \text{for all } \alpha, \beta \subseteq Z_n, \quad \alpha \cap \beta = \phi,$$

are special cases of (2). For related results see Carlson [2] and Fan [3], [4].

It is obvious that positive semi-definite and totally non-negative matrices are WSS, and it is easy to show that M -matrices (which we allow to have non-negative principal minors) are also WSS. It is also true that all matrices in these classes are GH. However, not all WSS-matrices are GH; the most obvious examples being the $k \times k$ cyclic permutation matrices with $k \geq 3$ and k odd.

THEOREM 1. *Every GH-matrix is a WSS-matrix. The WSS-matrix A is a GH-matrix if and only if*

$$\text{for } \alpha \subseteq \beta \subseteq Z_n, \quad A(\beta) = 0 \quad \text{whenever} \quad A(\alpha) = 0. \quad (3)$$

Proof. Gantmacher and Krein [6, p. 111] and Carlson [1] have shown that if all principal minors of A are positive, then A is GH if and only if it is WSS. Carlson's proof that GH-matrices are WSS actually holds in general. To prove our second statement, suppose first that A is GH. If $A(\alpha) = 0$ and $\alpha \subseteq \beta$, then

$$0 \leq A(\beta) \leq A(\alpha)A(\beta - \alpha) = 0,$$

and $A(\beta) = 0$. Conversely, suppose A is WSS and that (3) holds. Given $\alpha, \beta \subseteq Z_n$, either $A(\alpha \cup \beta) = 0$ (in which case $A(\alpha \cup \beta)A(\alpha \cap \beta) \leq A(\alpha)A(\beta)$, since $A(\alpha) \geq 0$, $A(\beta) \geq 0$) or $A(\alpha \cup \beta) > 0$ (in which case all principal minors of $A[\alpha \cup \beta]$ are positive, by (3), and Carlson's proof of (2) holds.

THE RANK OF A WSS-MATRIX

We next study the relationship between nested principal minors of general WSS-matrices. Let $\rho(A)$ denote the rank of A .

LEMMA 1. *Let A be an $n \times n$ WSS-matrix. Suppose $A(\alpha) > 0$ for some $\alpha \subseteq Z_n$, $0 \leq |\alpha| \leq n-2$, and $A(\alpha, s) = 0$ for some $s \in Z_n - \alpha$. For any $t \in Z_n - \{\alpha, s\}$, let $\gamma = \{\alpha, s, t\}$; then*

$$A(\gamma) = A(\alpha, s, t) = 0, \quad (4)$$

and

$$A(\alpha, s|\alpha, t)A(\alpha, t|\alpha, s) = 0. \quad (5)$$

Further, if $A(\alpha, s|\alpha, t) = 0$, then

$$A(\alpha, s|\gamma - r) = 0 \quad \text{for all } r \in \gamma. \quad (6)$$

If $A(\alpha, t|\alpha, s) = 0$ then

$$A(\gamma - r|\alpha, s) = 0 \quad \text{for all } r \in \gamma. \quad (6')$$

Proof. Let $d_{ij} = A(\alpha, i|\alpha, j)$ for $i, j \in \{s, t\}$. The hypotheses, together with Sylvester's identity (cf. [5, Vol. I, p. 33]) imply that

$$0 \leq A(\alpha)A(\gamma) = \det(d_{ij}) = -d_{st}d_{ts} = -A(\alpha, s|\alpha, t)A(\alpha, t|\alpha, s). \quad (7)$$

Since A is WSS, we must have equality in (7), which is (5); also, since $A(\alpha) > 0$, we have (4).

If $\alpha = \phi$, (6) and (6') are obvious. Assume then that $\alpha \neq \phi$. Suppose $A(\alpha, s|\alpha, t) = 0$. By hypothesis, since $A(\alpha) > 0$, the set C of columns of $A[\alpha, s|\alpha]$ is linearly independent. As $A(\alpha, s) = A(\alpha, s|\alpha, t) = 0$, the columns $A[\alpha, s|s]$ and $A[\alpha, s|t]$ are linearly dependent on C . Since all columns of $A[\alpha, s|\gamma]$ are linearly dependent on C , which has only $|\alpha|$ elements, (6) holds. If $A(\alpha, t|\alpha, s) = 0$, a similar argument gives (6').

Our next result is essentially known for positive semi-definite matrices. Since positive, semi-definite matrices are WSS, this follows from Lemma 1 above and Theorem 6 of [11, p. 92]. For related results for totally non-negative matrices, see Gantmacher and Krein [6, pp. 113–115] or Karlin [8, pp. 89–91].

THEOREM 2. *Let A be an $n \times n$ WSS-matrix. Suppose that for some p , $1 \leq p \leq n - 2$, that all principal minors of order p are positive, and all principal minors of order $p + 1$ are zero. Then the rank of A is p .*

Proof. We shall show that for every $\beta = \{\alpha, i\}$, $|\alpha| = p$, $i \notin \alpha$, and $j \notin \alpha$, the column $A[\beta|j]$ is linearly dependent on the columns of $A[\beta|\alpha]$. It will follow that for every β' , $\beta \neq \beta'$ and $|\beta'| = p + 1$, $A(\beta|\beta') = 0$. From this it will follow that $\rho(A) = p$.

Without loss of generality, we may assume $\alpha = \{1, 2, \dots, n - 2\}$, $i = n - 1$, $j = n$. Now all principal minors of order $n - 2$ are positive, and those of order $n - 1$ are zero. By Lemma 1, $A(n \supset 1|\hat{n})A(\hat{n}|n \supset 1) = 0$. By repeated application of Lemma 1, if $A(n \supset 1|\hat{n}) = 0$, then for every $k \in \alpha$, $A(\hat{k}|\hat{n}) = 0$, and hence $A(\hat{k}|n \supset 1) = 0$, and hence $A(\hat{n}|n \supset 1) = 0$. Now we have that $A(\hat{n}|n \supset 1) = 0$, so by the proof of Lemma 1, for $j \neq n$, $A[\hat{n}|j]$ is linearly dependent on the columns of $A[\hat{n}|\alpha]$.

COROLLARY. *Let A be an $n \times n$ WSS-matrix. Suppose for some p , $1 \leq p < n$, that all principal submatrices of order p are non-singular and generalized Hadamard and that all principal minors of order $p + 1$ are zero. Then A is a GH-matrix.*

Proof. The proof follows from Theorems 1 and 2.

A CHARACTERIZATION OF WSS-MATRICES WHICH ARE NOT GH

In this section we show that every WSS-matrix which is not GH is related to the example, given earlier, of a $k \times k$ cyclic permutation matrix.

LEMMA 2. *Let A be a WSS-matrix with $\det A > 0$ and $A[\alpha]$ a GH-matrix for all $\alpha \subsetneq Z_n$. The following are equivalent:*

- (a) *A is not a GH-matrix.*
- (b) *There exists $\alpha \neq Z_n$ such that $A(\alpha) = 0$.*
- (c) *$\text{Adj } A$ has a non-singular principal submatrix, of order $k \geq 3$, which is a cyclic permutation of a diagonal matrix.*

Proof. Conditions (a) and (b) are equivalent by Theorem 1.

Suppose (b) holds. We will first show that all principal minors of A of orders $n - 2$ and $n - 1$ are zero. By Theorem 1 there exists $\gamma_0 \subseteq Z_n$,

with $\alpha \subseteq \gamma_0$ and $|\gamma_0| = n - 1$, such that $A(\gamma_0) = 0$. Let $\gamma \subseteq Z_n$, $|\gamma| = n - 1$, with $\gamma \neq \gamma_0$. Let $\{i_0\} = Z_n - \gamma_0$ and $\{j_0\} = Z_n - \gamma$. Then $\gamma_0 \cap \gamma$ contains $n - 2$ elements of Z_n and $\gamma_0 \cap \gamma \subseteq \gamma_0$. Since $\det A > 0$, all principal minors of order $n - 2$ of $A[\gamma_0]$ are zero by Lemma 1. Therefore $A(\gamma_0 \cap \gamma) = 0$; and since $\gamma_0 \cap \gamma \subset \gamma$ and $A[\gamma]$ is a GH-matrix, $A(\gamma) = 0$. Thus all principal minors of order $n - 1$ of A are zero. It follows from this and $\det A > 0$, by Lemma 1, that all principal minors of order $n - 2$ of A are zero.

Thus by Jacobi's Theorem [5, Vol. I, p. 21] all principal minors of $B = \text{Adj } A$ are non-negative, and all diagonal elements and 2×2 principal minors of B are zero. Let k be the minimum value, $3 \leq k \leq n$, such that there exists $B[\psi]$ of order k with $B(\psi) > 0$ and $B(\alpha) = 0$ for all $\phi \neq \alpha \not\subseteq \psi$. Then by Theorem 2.1.10 of [7], $B[\psi]$ is a cyclic permutation of a non-singular diagonal matrix.

Conversely, suppose (c) holds. Since $B[\psi]$ and therefore $B = \text{Adj } A$ has a zero diagonal element, Jacobi's Theorem implies that A has a zero principal minor of order $n - 1$; and since $\det A > 0$, (b) holds.

For $1 \leq m \leq n$, let $C_m(A)$ denote the m th compound of A [10, p. 16]. Since by Theorem 1, A is a GH-matrix if and only if all principal submatrices of A are GH-matrices, the following holds.

THEOREM 3. *Let A be a WSS-matrix. A is not a GH-matrix if and only if for some m , $1 \leq m \leq n$, $C_{m-1}(A)$ has a principal submatrix of order $k \geq 3$, which is a cyclic permutation of a non-singular diagonal matrix, and which is determined by precisely m rows of A .*

The author wishes to thank Professor Carlson who has kindly shared his expertise and knowledge of matrix theory in the selection of relevant results to be included in this paper. His suggestions and contributions concerning notation and style have greatly improved the clarity of the paper.

REFERENCES

- 1 D. Carlson, Weakly sign-symmetric matrices and some determinantal inequalities, *Colloq. Math.* **27**(1967), 123-129.
- 2 D. Carlson, On some determinantal inequalities, *Proc. A. M. S.* **19**(1968), 462-466.
- 3 Ky Fan, An inequality for subadditive functions on a distributive lattice, with applications to determinantal inequalities, *Lin. Alg. Appl.* **1**(1968), 33-38.
- 4 Ky Fan, Subadditive functions on a distributive lattice and on extensions of Szasz's inequality, *J. Math. Anal. Appl.* **18**(1967), 262-268.
- 5 F. R. Gantmacher, *The Theory of Matrices*, Chelsea, New York (1960), 2 vols.

- 6 F. R. Gantmacher and M. G. Krein, *Oszillationsmatrizen, Oszillationskerne und kleine Schwingungen mechanischer Systeme*, Akademie-Verlag, Berlin (1960).
- 7 B. M. Green, Characterizations of matrices for which certain determinantal equalities hold, Ph.D. thesis, Oregon State University (1969), 75 numb. leaves (microfilm).
- 8 S. Karlin, *Total Positivity*, Stanford University Press, Stanford (1968).
- 9 D. M. Koteljanskii, A property of sign-symmetric matrices (Russian), *Ukrain. Mat. Z.* **2**(1950); English transl.: Ser. 2, **27**(1963), 19–23.
- 10 M. Marcus and H. Minc, *A Survey of Matrix Theory and Matrix Inequalities*, Allyn and Bacon, Boston (1964).
- 11 H. W. Turnbull and A. C. Aitken, *An Introduction to the Theory of Canonical Matrices*, Dover, New York (1961).

Received November 16, 1972